

# Rehydrat® Tabs

agilan® complementary feed

To support digestion  
highly soluble, effervescent, rich in energy and electrolytes

For calves, piglets and poultry

## Composition

Lactose; Sodium bicarbonate; Sodium chloride; Potassium chloride

## Analytical constituents

Crude protein	5.5 %
Crude fat	1.8 %
Crude fiber	0.5 %
Crude ash	17.9 %
Sodium	6.5 %
Potassium	1.2 %
Chloride	4.3 %
Lysine	0.0 %
Methionine	0.0 %

## Content of additives per kg

<u>Preservatives:</u>	Citric acid (E330)	80,000 mg
<u>Vitamins:</u>	Vitamin C (3a300)	10,000 mg

## Recommended use

For oral rehydration in *calves, piglets and poultry*.

Rehydrat® Tabs quickly and reliably compensate electrolyte and body water losses due to an adequate mineral compositions and energy deficiencies due to an energy-rich lactose.

## Feeding recommendation

*Calves:* 1 tab/liter of lukewarm water

should be given additionally to the usual milk meals as 1 -3 snacks per day in a daily amount corresponding to the severity of the cases for independent intake.

The Rehydrat® solution should be used for the entire duration of an altered fecal consistency.

Fecal consistency	Additional liquid requirement (in % of b.w.)	40 kg calf	
		per day	waterings per day
Thin mushy	approx. 5 %	2 l	2 l / 1 x daily
Soupy	approx. 10 %	4 l	2 l / 2 x daily
Watery	15 (-20) %	6 (-8 l)	2 l / 3 x daily

*Piglets:* 1 tab/liter of lukewarm water

The Rehydrat® solution is offered to the piglets for autonomous intake. It is favourable to prepare the solution fresh several times a day. This maintains the piglets' curiosity and prevents odor absorption from the environment.

*Poultry:* Dissolve 1 tab/100 liters of drinking water.



# Rehydrat® Tabs

## Feeding instruction

The solution should be freshly prepared with clean water prior to each use.

Rehydrat® Tabs should be administered at intervals to the ration of whole milk / milk replacer. Due to lactose Rehydrat® Tabs are very rich in energy with a low osmolarity. Lactose maintains the lactase activity so that it is possible to return to milk feeding without any problems.

This product must not be used as a sole drinker for calves and piglets.

## Recommended feeding period: 1 - 7 days

In case of risk of, during periods of or after digestive disturbance.

It is recommended that a veterinarian's opinion be sought before use or before extending the feeding period.

## Storage

Shelf life guarantee is only valid for unopened packages, stored at a cool and dry place.

Keep the product out of the reach of children!

Status: February 2019